

TEST: Progesterone

PRINCIPLE:

Progesterone (4-pregnen-3.20-dione) is one of the 21 carbon steroids secreted by the corpus luteum of the ovary in females during the normal menstrual cycle. It is also produced in low concentrations by the adrenal cortex in both males and females. In pregnancy the placenta is the major source of progesterone after the seventh gestational week. Progesterone has two main biological functions. First, it transforms the estrogen stimulated endometrium into the secretory phase, which allows implantation of the fertilized ovum. Secondly, it sustains the pregnancy by decreasing uterine contractility. During the follicular phase progesterone concentrations are low prior to the mid-cycle gonadotropin surge. Progesterone concentrations in the range 0.3-3.5 nmol/L have been measured the day before the LH peak. Immediately after the LH surge, concentrations begin to rise rapidly and reach maximum levels at the middle of the luteal phase. During a normal luteal phase progesterone concentration are between 3.9-80 nmol/L and the peak level should exceed 25 nmol/L.Measurements of maternal progesterone levels have been suggested for the clinical assessment of threatening abortion, hydatidiform mole and rhesus isoimmunisation.

SPECIMEN REQUIREMENTS:

2ml collected in a serum separator tube (gel barrier). Separate serum from cells ASAP or within 2 hours of collection by centrifugation. Stability after separation from cells: Ambient: 8 Hours; Refrigerated: 48 Hours; Frozen: 1 year (avoid repeated freeze/thaw cycles).

REJECTION CRITERIA: Plasma or other body fluids. Gross hemolysis

METHOD: Enhanced Chemiluminescence.

REFERENCES:

1. Cooke, I.D. (1976): In Loraine, J.A. and Bell, E.T. (eds): Hormone assays and their clinical application. Churchill Livingstone, Edinburgh, London and New York, 4th edition, pp 447-518.

Normal Range: Female:

Mid Follicular Phase 0.31-1.52 ng/ml
Mid-Luteal Phase 5.16-18.56 ng/ml
Post Menopausal Less than 0.7 ng/ml

Pregnancy:

First Trimester 4.73 – 50.74 ng/ml Second Trimester 19.41 – 445.30 ng/ml

Male 0.14-2.06 ng/ml

Turnaround Time: 2 business days